

Cáncer rectal

Bibliografía

1. Office of National Statistics, Cancer Statistics registrations: Registrations of cancer diagnosed in 2004, England. Series MB1 no; 35
2. Guidelines for the management of colorectal cancer (2001). Association of Coloproctology of Great Britain and Ireland
3. McSherry CK, Cornell GN, Glenn F. Carcinoma of the colon and rectum. *Ann Surg* 1969; 169: 502-09
4. Dixon AR, Thornton-Holmes J, Cheetham NM. General practitioners' awareness of colorectal cancer: a 10 year review. *BMJ* 1990; 301: 445-46
5. Bulmer M, Hartley J, Lee PW, Duthie GS et al. Improving the view in the rectal clinic: a randomized control trial. *Ann R Coll Surg Engl* 2000; 82: 210-12
6. Farouk R, Edwards J, Thorne M et al. Brush cytology for the diagnosis of rectal carcinoma. *Br J Surg* 1996; 83: 1456-58
7. Schaffzin DM, Wong WD. Endorectal ultrasound in the preoperative evaluation of rectal cancer. *Clin Colorectal Cancer* 2004; 4: 124-32
8. Solomon MJ, McLeod RS. Endoluminal transrectal ultrasonography: accuracy, reliability, and validity. *Dis Colon Rectum* 1993; 36: 200-05
9. Akasu T, Sugihara K, Moriya Y, Fujita S. Limitations and pitfalls of transrectal ultrasonography for staging of rectal cancer. *Dis Colon Rectum* 1997; 40(suppl): S10-S15.
10. Garcia-Aguilar J, Pollack J, Lee SH et al. Accuracy of endorectal ultrasonography in preoperative staging of rectal tumors. *Dis Colon Rectum* 2002; 45: 10-15
11. Zammit M, Jenkins JT, Urie A et al. A technically difficult endorectal ultrasound is more likely to be inaccurate. *Colorectal Dis* 2005; 7: 486-91
12. Kwok H, Bissett IP, Hill GL. Preoperative staging of rectal cancer. *Int J Colorectal Dis* 2000; 15: 9-20
13. Guinet C, Buy JN, Ghossain MA et al. Comparison of magnetic resonance imaging and computer tomography in the preoperative staging of rectal cancer. *Arch Surg* 1990; 125: 385-88
14. Butch RJ, Stark DD, Wittenberg J et al. Staging rectal cancer by MR and CT. *AJR Am J Roentgenol* 1986; 146: 1155-60
15. Indinnimeo M, Grasso RF, Cicchini C et al. Endorectal magnetic resonance imaging in the preoperative staging of rectal tumours. *Int Surg* 1996; 81: 419-22
16. Birbeck K, Macklin C, Tiffin NJ et al. Rates of circumferential resection margin involvement vary between surgeons and predict outcomes in rectal cancer surgery. *Ann Surg* 2002; 4: 449-57
17. Adam IJ, Mohamdee MO, Martin IG et al. Role of circumferential margin involvement in the local recurrence of rectal cancer. *Lancet* 1994; 344: 707-11
18. Beets-Tan RG, Beets GL, Vliegen RF et al. Accuracy of magnetic resonance imaging in prediction of tumor-free resection margin in rectal cancer surgery. *Lancet* 2001; 357: 497-504
19. Chen CC, Lee RC, Lin JK et al. How accurate is magnetic resonance imaging in restaging rectal cancer in patients receiving preoperative combined chemoradiotherapy? *Dis Colon Rectum* 2005; 48: 722-28
20. Kapiteijn E, Marijnen C, Colenbrander AC et al. Local recurrence in patients with rectal cancer, diagnosed between 1988 and 1992: a population-based study in the west Netherlands. *Eur J Surg Oncol* 1998; 24: 528-35
21. Heald RJ, Moran BJ, Ryall RD et al. Rectal cancer: the Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998; 133: 894-99

22. Glimelius B, Isacsson U, Jung B et al. Radiotherapy in addition to radical surgery in rectal cancer-evidence for a dose-response effect favouring preoperative treatment. *Int J Radiat Oncol Biol Phys* 1997; 15: 281-87
23. Minsky BD, Cohen AM, Enker WE et al. Combined modality therapy of rectal cancer: decreased acute toxicity with the pre-operative approach. *J Clin Oncol* 1992; 10: 1218-24
24. Janjan NA, Khoo VS, Abbruzzese J et al. Tumor downstaging and sphincter preservation with preoperative chemoradiation in locally advanced rectal cancer: the M.D. Anderson Cancer Center experience. *Int J Radiat Oncol Biol Phys* 1999; 44: 1027-38
25. Improved survival with preoperative radiotherapy in resectable rectal cancer. Swedish Rectal Cancer Trial. *N Eng J Med* 1997; 336:980-87
26. Kapiteijn E, Marijnen CA, Nagtegaal ID et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Eng J Med* 2001; 345:638-46
27. Quirke P, Sebag-Montefiore D, Steele R et al. Local recurrence after rectal cancer resection is strongly related to the plane of surgical dissection and is further reduced by pre-operative short course radiotherapy. Preliminary results of the Medical Research Council (MRC) CR07 trial. *Journal of Clinical Oncology*, 2006 ASCO Annual Meeting Proceedings Part I. Vol 24, No. 18S (June 20 Supplement), 2006: 3512
28. Sauer R, Becker H, Hohenberger W et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Eng J Med* 2004; 351:1731-40
29. Pahlman L, Glimelius B. Pre or postoperative radiotherapy in rectal and rectosigmoid carcinoma. Report from a randomized multicenter trial. *Ann Surg* 1990; 211: 187-95
30. Hyams DM, Mamounas EP, Petrelli N et al. A clinical trial to evaluate the worth of preoperative multimodality therapy in patients with operable carcinoma of the rectum: a progress report of National Surgical Adjuvant Breast and Bowel Project Protocol R-03. *Dis Colon Rectum* 1997; 40: 131-39
31. Guillem JG, Chessin DB, Cohen AM et al. Long-term oncologic outcome following preoperative combined modality therapy and total mesorectal excision of locally advanced rectal cancer. *Ann Surg* 2005; 241: 829-38
32. Gerard JP, Chapet O, Nemoz C et al. Preoperative concurrent chemoradiotherapy in locally advanced rectal cancer with high-dose radiation and oxaliplatin-containing regimen: the Lyon R0-04 phase II trial. *J Clin Oncol* 2003; 21: 1119-24
33. Dunst J, Reese T, Sutter T et al. Phase I trial evaluating concurrent combination of radiotherapy and capecitabine in rectal cancer. *J Clin Oncol* 2002; 20: 3983-91
34. Gastrointestinal Tumor Study Group. Prolongation of the disease-free interval in surgically treated rectal carcinoma. *N Eng J Med* 1985; 312: 1465-72
35. Fisher B, Wolmark N, Rockette H et al. Postoperative adjuvant chemotherapy or BCG for colon cancer: results from NSABP Prortocol R-01. *J Natl Cancer Inst* 1988; 80: 21-29
36. Krook JE, Moertel CG, Gunderson LL et al. Effective surgical adjuvant therapy for high-risk rectal cancer. *N Eng J Med* 1991; 324: 709-15
37. NIH consensus conference: adjuvant therapy for patients with colon and rectal cancer. *JAMA* 1990; 264: 1444-50
38. Wolmark N, Weiland HS, Hyams DM et al. Randomized trial of postoperative adjuvant chemotherapy with or without radiotherapy for carcinoma of the rectum: National Surgical Adjuvant Breast and Bowel Project Protocol R-02. *J Natl Cancer Inst* 2000; 92: 388-96
39. Baxter NN, Rothenberger DA, Morris AM et al. Adjuvant radiation for rectal cancer: Do we measure up to the standard of care? An epidemiologic analysis of trends over 25 years in the United States. *Dis Colon Rectum* 2005; 48: 9-15
40. Ortholan C, Francois E, Thomas O et al. Role of radiotherapy with surgery for T3 and respectable T4 rectal cancer: evidence from randomized trials. *Dis Colon Rectum* 2006; 49: 302-10
41. Matthiessen P, Hallbook O, Rutegard J et al. Population-based study of risk factors for postoperative death after anterior resection of the rectum. *Br J Surg* 2006; 93: 498-503

42. Enker WE, Merchant N, Cohen AM et al. Safety and efficacy of low anterior resection for rectal cancer, 681 consecutive cases from a specialty service. *Ann Surg* 1999; 230: 544-54
43. Williams N, Seow-Cohen F. Physiological and functional outcome following ultra-low anterior resection with colon-pouch anastomosis. *Br J Surg* 1998; 85: 1029-35
44. Nesbakken A, Nygaard K, Bull-Njaa T et al. Bladder and sexual dysfunction after mesorectal excision for rectal cancer. *Br J Surg* 2000; 87: 206-10
45. Whiteway J, Nicholls RJ, Morson BC. The role of surgical excision in the treatment of rectal cancer. *Br J Surg* 1985; 72: 694-97
46. Michelassi F, Vannucci L, Montag A et al. Importance of tumour morphology for the long term prognosis of rectal adenocarcinoma. *Am Surg* 1988; 54: 376-79
47. Chambers WM, Khan U, Gagliano A et al. Tumour morphology as a predictor of outcome after local excision of rectal cancer. *Br J Surg* 2004; 91: 457-59
48. Sitzler PJ, Seow-Choen F, Ho YH, et al. Lymph node involvement and tumor depth in rectal cancers: an analysis of 805 patients. *Dis Colon Rectum* 1997; 40: 1472-76
49. Brodsky JT, Richard GK, Cohen AM et al. Variables correlated with the risk of lymph node metastasis in early rectal cancer. *Cancer* 1992;69: 322-26
50. Bouvet M, Milas M, Giacco GG et al. Predictors of recurrence after local excision and postoperative chemoradiation therapy of adenocarcinoma of the rectum. *Ann Surg Oncol* 1999; 6: 26-32
51. Graham RA, Garnsey L, Jessup JM. Local excision of rectal cancer. *Am J Surg* 1990; 160: 306-12
52. Beets-Tan RG, Beets GL. Rectal cancer: review with emphasis on MR imaging. *Radiology* 2004; 232: 335-46
53. Zenni GC, Abraham K, Harford FJ et al. Characteristics of rectal carcinomas that predict the presence of lymph node metastases: implications for patient selection for local therapy. *J Surg Oncol* 1998; 67: 99-103
54. Blumberg D, Paty PB, Guillem JG et al. All patients with small intramural rectal cancers are at risk for lymph node metastasis. *Dis Colon Rectum* 1999; 42: 881-85
55. Minsky BD, Rich T, Recht A et al. Selection criteria for local excision with or without adjuvant radiation therapy for rectal cancer. *Cancer* 1989; 63: 1421-29
56. Consorti F, Lorenzotti A, Midiri G et al. Prognostic significance of mucinous carcinoma of colon and rectum: a prospective case-control study. *J Surg Oncol* 2000; 73: 70-74
57. Christiansen J. Excision of mid-rectal lesions by the Kraske sacral approach. *Br J Surg* 1980; 67: 651-52
58. Bergman L, Solhaug JH. Posterior trans-sphincteric resection for small tumours of the lower rectum. *Acta Chir Scand* 1986; 152: 313-16
59. Blair S, Ellenhorn JD. Transanal excision for low rectal cancers is curative in early- stage disease with favorable histology. *Am Surg* 2000; 66: 817-20
60. Chakravarti A, Compton CC, Shellito PC et al. Long-term follow-up of patients with rectal cancer managed by local excision with and without adjuvant irradiation. *Ann Surg* 1999; 230: 49-54
61. Weiser MR, Landmann RG, Wong WD et al. Surgical salvage of recurrent rectal cancer after transanal excision. *Dis Colon Rectum* 2005; 48: 1169-75
62. Hahnloser D, Wolff BG, Larson DW et al. Immediate radical resection after local excision of rectal cancer: an oncologic compromise? *Dis Colon Rectum* 2005; 48: 429-37
63. Buess G, Kipfmuller K, Ibaldo R et al. Clinical results of transanal endoscopic microsurgery. *Surg Endosc* 1988; 2: 245-50
64. Lee W, Lee D, Choi S et al. Transanal endoscopic microsurgery and radical surgery for T1 and T2 rectal cancer. *Surg Endosc* 2003; 17: 1283-87
65. Neary P, Makin GB, White TJ et al. Transanal endoscopic microsurgery: a viable alternative in selected patients with rectal lesions. *Ann Surg Oncol* 2003; 10: 1106-11

66. Winde G, Nottberg H, Keller R et al. Surgical cure for early rectal carcinomas (T1): transanal endoscopic microsurgery vs anterior resection. *Dis Colon Rectum* 1996; 39: 969-76
67. Heintz A, Morschel M, Junginger T. Comparison of results after transanal endoscopic microsurgery and radical resection for T1 carcinoma of the rectum. *Surg Endosc* 1998; 12: 1145-48
68. Moore HG, Guillem JG. Local therapy for rectal cancer. *Surg Clin N Am* 2002; 82: 967-81
69. Steele GD, Herndon JE, Bleday RR et al. Sphincter-sparing treatment for distal rectal adenocarcinoma. *Ann Surg Oncol* 1999; 6: 413-15
70. Russell AH, Harris J, Rosenberg PJ et al. Anal sphincter conservation for patients with adenocarcinoma of the distal rectum: long term results of radiation therapy oncology group protocol 89-02. *Int J Radiat Oncol Biol Phys* 2000; 46: 313-22
71. Perretta S, Guerrero V, Garcia-Aguilar J. Surgical treatment of rectal cancer: local resection. *Surg Oncol Clin N Am* 2006; 15: 67-93
72. Kim JS, Kim JS, Cho MJ et al. Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. *Int J Radiat Oncol Biol Phys* 2002 54: 403-08
73. Lezoche E, Guerrieri M, Paganini AM et al. Transanal endoscopic versus total mesorectal laparoscopic resections of T2-N0 low rectal cancers after neoadjuvant treatment: a prospective randomized trial with a 3-years minimum follow-up period. *Surg Endosc* 2005; 19: 751-56.