

Are Lifestyle Measures Effective in Patients With Gastroesophageal Reflux Disease?

An Evidence-Based Approach

Tonya Kaltenbach, MD; Seth Crockett, MD; Lauren B. Gerson, MD, MSc

Lifestyle modifications are first-line therapy for patients with gastroesophageal reflux disease (GERD). We applied an evidence-based approach to determine the efficacy of lifestyle measures for GERD management. We used PubMed and Ovid to perform a search of the literature published between 1975 and 2004 using the key words *heartburn, GERD, smoking, alcohol, obesity, weight loss, caffeine or coffee, citrus, chocolate, spicy food, head of bed elevation, and late-evening meal*. Each study was reviewed by 2 reviewers who assigned one of the following ratings: evidence A, randomized clinical trials; evidence B, cohort or case-control studies; evidence C, case reports or flawed clinical trials; evidence D, investigator experience; or evidence E, insufficient information. We screened 2039 studies and identified 100 that were relevant. Only 16 clinical trials examined the impact on GERD (by change in symptoms, esophageal pH variables, or lower esophageal sphincter pressure) of the lifestyle measure. Although there was physiologic evidence that exposure to tobacco, alcohol, chocolate, and high-fat meals decreases lower esophageal sphincter pressure, there was no published evidence of the efficacy of dietary measures. Neither tobacco nor alcohol cessation was associated with improvement in esophageal pH profiles or symptoms (evidence B). Head of bed elevation and left lateral decubitus position improved the overall time that the esophageal pH was less than 4.0 (evidence B). Weight loss improved pH profiles and symptoms (evidence B). Weight loss and head of bed elevation are effective lifestyle interventions for GERD. There is no evidence supporting an improvement in GERD measures after cessation of tobacco, alcohol, or other dietary interventions.

Arch Intern Med. 2006;166:965-971

Gastroesophageal reflux disease (GERD) is characterized by symptoms of substernal burning or acid regurgitation produced by the abnormal reflux of gastric contents into the esophagus.¹ According to US population surveys, 44% of all Americans experience heartburn at least once per month, 14% at least once per week, and up to 7% daily.^{2,3} The major mechanism for GERD is transient relaxation of the lower esophageal sphincter (LES).⁴ Recommendations for lifestyle modifications are based on the presumption that certain foods, body position, tobacco, alcohol, and obesity contribute to

a dysfunction in the body's antireflux defense system. Accordingly, the American College of Gastroenterology recommends the use of lifestyle changes, including elevation of the head of the bed (HOB); decreased fat, chocolate, alcohol, peppermint, and coffee intake; cessation of smoking; and avoiding recumbency for 3 hours postprandially, in addition to antireflux medical treatment.¹ However, the evidence

*CME course available at
www.archinternmed.com*

to support such lifestyle modification recommendations has not been well substantiated. The purpose of this study is to apply an evidence-based approach to

Author Affiliations: Division of Gastroenterology and Hepatology (Drs Kaltenbach and Gerson) and Department of Medicine (Dr Crockett), Stanford University School of Medicine, Stanford, Calif.

Table 1. Evidence Levels

Evidence A	Evidence B	Evidence C	Evidence D	Evidence E
One or more well-designed randomized controlled clinical trials with consistent evidence	Cohort or case-control trials, nonrandomized or uncontrolled clinical trials	Case reports, flawed clinical trials, population studies	Expert or investigator opinion	Insufficient evidence or trials with significantly conflicting data

Table 2. Summary of Physiologic Evidence*

Factor	Trials, No.	Lowered LESP	Worsened pH	Worsened Symptoms	References
Tobacco	12	B	B	B	6-17
Alcohol	16	No effect (B)	B	B	3,6,9,10,16-27
Obesity	24	E	E	E	28-51
Coffee and caffeine	14	E	E	No effect (C)	10,16,17,52-62
Chocolate	2	B	B	E	63,64
Spicy foods	2	E	E	C	3,65
Citrus	3	No effect (B)	E	C	52,66,67
Carbonated beverages	2	B	E	C	68,69
Fatty foods	9	D	B	E	70-78
Mint	1	D	E	E	79
Recumbent position	1	E	B	B	80
RLD position	3	B	B	E	81-83
Late-evening meal	3	E	No effect (B)	E	84-86

Abbreviations: LESP, lower esophageal sphincter pressure; RLD, right lateral decubitus.

*See Table 1 for a summary of evidence A through E.

the literature to determine which lifestyle measures are most likely to benefit patients with GERD.

METHODS

We performed a systematic review of the English-language literature relating to human subjects published between 1975 and 2004 using PubMed and Ovid to obtain data pertaining to the effects of lifestyle modifications on GERD. We included the following key words in the search: *lifestyle modification, heartburn, GERD, reflux, smoking, alcohol, obesity, weight loss, caffeine or coffee, citrus, chocolate, mint, carbonated beverages, spicy food, fatty foods, head of bed elevation, and late-evening meal.*

To be included in the study, trials had to contain a lifestyle intervention and outcomes of GERD measures, including heartburn symptoms, ambulatory esophageal pH monitoring, and esophageal manometric variables. Studies of the physiologic effects of lifestyle interventions only are included as introductory evidence for each lifestyle intervention.

We applied an evidence-based approach to the literature using a standard scoring system (Table 1).⁵ After 2 of us (either T.K. and L.B.G. or S.C. and L.B.G.) independently reviewed each

study, evidence-based ratings of A through E were assigned. A consensus was obtained through a second review if there was a difference of opinion on initial review.

RESULTS

Of 2039 trials screened for relevancy, 100 were included in this analysis. Only 16 trials examined the impact of lifestyle changes on GERD variables. The subsections that follow describe the physiologic evidence of lifestyle interventions on GERD measures (Table 2), followed by the available evidence on cessation of the lifestyle measure and the impact on GERD variables (Table 3).

SMOKING

Cigarette smoking has been associated with symptomatic GERD. Questionnaires have reported higher rates of reflux symptoms in tobacco smokers compared with nonsmokers.⁶⁻⁹ Multivariate analysis from a case-control study¹⁰ showed that duration of smoking was associated with

increasing reflux symptoms, with an odds ratio (OR) for reflux of 1.7 (95% confidence interval [CI], 1.4-2.0; $P < .001$) in daily smokers with a greater than 20-year tobacco use history compared with those who smoked daily for less than 1 year. Cigarette smoking has been shown to prolong acid clearance and decrease baseline LES pressure (LESP),^{96,97} with an abrupt decline in LESP at the start of smoking that returns to normal within minutes of completion of the cigarette.⁹⁸ Abrupt increases in intra-abdominal pressure, such as with coughing or deep inspiration, have been associated with reflux symptoms in smokers.¹¹ Studies¹²⁻¹⁴ that examined acid perfusion (using the Bernstein test) or esophageal pH have reported conflicting associations; for example, despite having more "reflux episodes," smokers (compared with nonsmokers) do not show increased esophageal acid exposure time.¹⁵

No improvement was shown in 3 case-control studies (evidence B) that evaluated the effect of tobacco cessation on GERD outcomes. One study¹⁵ compared a cohort of 30 healthy volunteers (15 smokers and 15 nonsmokers) with 10 smokers with GERD and found that smokers had more reflux episodes than nonsmokers but that 24-hour smoking cessation did not reduce the total time that the esophageal pH was less than 4.0. Another study⁸⁷ showed that 1-day cessation of smoking decreased the number of daily reflux episodes but did not significantly affect total esophageal acid exposure. On the contrary, Kadakia et al,⁸⁸ after their evaluation of smokers with GERD who abstained from smoking for 48 hours, concluded that smoking significantly increased distal esophageal acid exposure. Although tobacco use has been associated with an adverse effect on the LES, the evidence to date does not support an improvement in GERD after cessation of tobacco use.

ALCOHOL

Alcohol consumption may precipitate GERD by increasing acid secretion through gastrin stimulation, reducing LESP, increasing spontaneous LES relaxations, and impairing

esophageal motility and gastric emptying.¹⁸ Modest alcohol intake has been shown to induce reflux symptoms and decrease esophageal pH in healthy individuals without GERD symptoms¹⁹⁻²¹ despite normal overall 24-hour pH measurements.²² Randomized and cross-sectional studies^{16,23-26} have suggested an increased prevalence of symptomatic reflux in alcohol users. Wang et al¹⁶ reported reflux symptoms in 43% of heavy (≥ 210 g/wk) alcohol users compared with 16% of nondrinkers (OR, 2.85; 95% CI, 1.67-4.49; $P < .01$); however, large American⁹ and multinational¹⁷ cross-sectional studies have not shown similar associations.

In a case-control study²⁷ (evidence B) that examined GERD outcomes in symptomatic alcoholic patients compared with matched groups (without alcohol consumption) of controls, patients with GERD, and patients with nutcracker esophagus, most alcoholic patients demonstrated LES hypertension, high-amplitude esophageal contractions, or nonperistaltic esophageal contractions that improved after prolonged (> 6 months) alcohol abstinence but were not accompanied by improvements in esophageal pH. Therefore, although esophageal motility abnormalities were shown to improve after the cessation of alcohol use, there is insufficient evidence to support the direct effect of alcohol abstinence on pH or GERD symptoms.

OBESITY

Obesity has been speculated to cause GERD symptoms because of multiple factors, including increased (1) gastroesophageal sphincter gradient,⁹⁹ (2) incidence of hiatal hernia,¹⁰⁰⁻¹⁰² (3) intra-abdominal pressure,¹⁰³ and (4) output of bile and pancreatic enzymes.¹⁰⁴ Although some studies²⁸ have shown increased GERD in morbidly obese individuals, other studies^{29,30} did not demonstrate similar findings. Several population-based studies³¹⁻³³ have found a significant relationship between increasing body mass index (calculated as weight in kilograms divided by the square of height in meters) and GERD symptoms. Murray

Table 3. Summary of Lifestyle Intervention or Cessation Trials*

Factor	Trials, No.	Raised LESP	Improved pH	Improved Symptoms	References
Tobacco	3	E	No effect (B)	No effect (B)	15,87,88
Alcohol	1	No effect (B)	No effect (B)	NA	27
Weight loss	5	E	B	B	89-93
Coffee and caffeine	0	E	E	E	NA
Chocolate	0	E	E	E	NA
Spicy foods	0	E	E	E	NA
Citrus	0	E	E	E	NA
Carbonated beverages	0	E	E	E	NA
Fatty foods	0	E	E	E	NA
Mint	0	E	E	E	NA
Elevation of the HOB	3	E	A	B	80,94,95
LLD position	3	B	B	E	81-83
Late-evening meal	2	E	B	E	84,85

Abbreviations: HOB, head of the bed; LESP, lower esophageal sphincter pressure; LLD, left lateral decubitus; NA, not applicable.

*See Table 1 for a summary of evidence A through E.

et al³¹ reported an adjusted OR of 1.82 (95% CI, 1.33-2.5) for overweight patients with weekly heartburn symptoms compared with 1.5 (95% CI, 1.13-1.99) for individuals of average weight. Cross-sectional studies³⁴⁻³⁸ have similarly reported higher weekly reflux symptoms in obese patients. Another study³⁹ showed that for every increment of body mass index of 5, the risk of GERD increased by 1.2. In a multivariate logistic regression model,³² women with a body mass index greater than 35 were shown to have an increased risk of GERD (OR, 6.3; 95% CI, 2.4-4.7) compared with men with a similar body mass index (OR, 3.3; 95% CI, 2.4-4.7; $P < .001$). Although some studies⁴⁰⁻⁴⁵ have reported a link between obesity and GERD by demonstrating abnormal pH and manometric measures, other studies⁴⁶⁻⁵¹ have refuted such an association.

Bariatric surgery has been associated with improved GERD variables.¹⁰⁵⁻¹¹² Given that the surgical procedure has inherent antireflux properties, it is difficult to directly assess the effect of weight loss on GERD. Five studies⁸⁹⁻⁹³ have evaluated the independent effect of weight loss on GERD variables. Fraser-Moodie et al⁸⁹ showed a significant correlation between weight loss and esophageal pH (OR, 0.55; $P < .001$) in an uncontrolled study of 34 obese patients with GERD. Mathus-Vliegen et al⁹¹⁻⁹³ demonstrated a similar correlation with a decreased upright pH less than 4 (8.0% vs 5.5%;

$P < .05$) and postprandial reflux episodes (49.0 vs 32.1; $P < .05$) in patients with a mean weight loss of 12.4 kg in 13 weeks. However, in another study⁹⁰ that randomized 20 obese patients with reflux esophagitis to either a 430-kcal/d diet or an unrestricted diet, there was no significant difference in reflux symptoms between controls and patients with a 10% body weight loss after 6 months. Therefore, although weight loss seems to have a promising effect on pH measures and symptoms, further randomized controlled studies are warranted to determine its exact effect on GERD outcomes.

CITRUS FRUITS AND JUICES

Citrus fruits are often cited as precipitants of heartburn symptoms. In a questionnaire of approximately 400 patients with GERD, 72% of respondents reported increased heartburn with ingestion of either orange or grapefruit juice.⁶⁶ However, Price et al⁵² found that acid-sensitive patients (positive Bernstein test results) were sensitive to intraesophageal infusion of orange juice, even when the orange juice was adjusted to a pH of 7, suggesting that acidity is not the only factor determining the citrus effect on GERD. Cranley et al⁶⁷ demonstrated that patients with GERD did not change their LESP when challenged with orange juice infusion; this was in contrast to controls who showed increased LESP. No stud-

ies, to our knowledge, have assessed the effect of avoidance of citrus on GERD.

CARBONATED BEVERAGES

Studies have postulated that carbonated beverages cause GERD.¹¹³ In one study,⁶⁸ carbonated soft drink consumption was found to be a predictor of GERD symptoms in a multivariate analysis. Another small study⁶⁹ of healthy individuals found equivalent decreases in LESP with ingestion of carbonated water, caffeinated cola, or caffeine-free cola compared with water ingestion. No cessation trials have been published to date, to our knowledge.

COFFEE AND CAFFEINE

Despite the fact that intraesophageal infusion of coffee in patients with acid sensitivity has been shown to cause heartburn,⁵² 2 large epidemiologic studies^{16,17} found no association between coffee drinking and GERD. A recent survey¹⁰ in Norway described a negative association between coffee drinking and GERD incidence. Although some studies^{53,54} have described increased LESP with regular and decaffeinated coffees, another trial⁵⁵ showed no effect of coffee on the number of reflux episodes, total reflux time, or LESP in patients with GERD or controls. There is conflicting evidence regarding how different coffee preparations, roasting methods, and processing methods affect GERD variables.⁵⁶⁻⁵⁹ Decaffeinated compared with caffeinated coffee ingestion was associated with significantly less acid exposure time in some studies.⁶⁰ Salmon et al⁶¹ noted that patients had lower postprandial LESP regardless of whether they were exposed to coffee, suggesting that the association between coffee and GERD may be confounded by the fact that people frequently drink coffee after meals. Another study⁶² reported that coffee decreased LESP in healthy individuals and in patients with reflux esophagitis when fasting or following a standard test meal. Given the conflicting reported data, the relationship between caffeine or coffee and GERD remains unclear. There

is insufficient evidence to support the routine recommendation that patients with GERD avoid such beverages.

CHOCOLATE

Chocolate is frequently implicated as a provoker of GERD. There are limited data that chocolate may affect esophageal pH and LES profiles. One early uncontrolled study⁶³ found that ingestion of 120 mL of chocolate syrup by 9 controls caused LESP to significantly decrease compared with baseline measurements ($P < .01$). In 7 patients with typical GERD symptoms, Murphy and Castell⁶⁴ found greater acid exposure time after chocolate ingestion compared with ingestion of a test drink of equivalent osmolality and caloric value ($P = .04$). No studies have addressed the effect of chocolate abstinence on GERD symptoms.

SPICY FOODS

Although patients frequently cite spicy foods as an inciting agent in heartburn episodes, little data have been published regarding the effects of spicy food consumption on LESP or esophageal pH. Nebel et al³ conducted a survey of patients with GERD symptoms and found that 88% listed spicy foods as a precipitant of heartburn. One study⁶⁵ found that in patients with GERD vs controls, onion intake increased the number of reflux episodes ($P < .01$) and the esophageal acid exposure time ($P < .05$). The results of this study have not been confirmed by other larger trials. Many patients who believe that spicy foods exacerbate their GERD symptoms may abstain from these foods although we found no published trials that studied the effect of abstinence on change in GERD symptoms.

FATTY FOODS

It has been postulated that fat increases not only reflux but also the sensitivity of the esophagus to acid exposure. In a randomized clinical trial,⁷⁰ 8 patients with GERD and 11 controls received either an intragastric infusion of isotonic sodium chloride or a 20% lipid solution fol-

lowed by an intraesophageal infusion of hydrochloric acid. There were no significant differences in acid sensitivity symptoms with respect to the fat infusion.⁷⁰ Two studies^{71,72} reported significantly shorter onset to heartburn symptoms or reflux episodes with acid infusion regardless of whether patients were receiving intraduodenal infusions of a lipid solution or isotonic sodium chloride. Fat infusions have not been shown to affect LESP or the rate of transient LES relaxations.⁷²

Nebel and Castell⁷³ evaluated the LES response to ingestion of fat and protein meals with equivalent caloric value in healthy individuals and found that fatty meals decreased LES pressure significantly compared with protein meals, which increased LESP. Becker et al⁷⁴ studied esophageal acid exposure after high- and low-fat meals and found that the fat content was not significantly associated with esophageal pH abnormalities. In a study⁷⁵ of 20 healthy individuals in the supine position, those who consumed high-fat meals had significantly increased acid exposure compared with the low-fat group, but this study also found that acid exposure increased in patients who ingested a greater volume. Other randomized trials have found no difference comparing high- and low-fat meals with respect to LESP, transient LES relaxations,⁷⁶ number of reflux episodes, or esophageal exposure to acid.⁷⁷ There is conflicting evidence regarding whether caloric content of foods is more important than fat content.^{76,78}

Based on the previously mentioned trials, there is limited evidence to suggest that the fat content of meals affects GERD outcomes. In fact, a recent review¹¹⁴ on this subject concluded that present evidence does not support any recommendations for patients with GERD to limit fat in their diets.

MINT

Peppermint and spearmint are plant extracts that are commonly used in foods and flavorings of toothpaste, mouthwash, and candy. Mint is commonly thought to relax the LES, and thus it is commonly recommended that patients with GERD

avoid ingesting mint.¹¹⁵ We found only 1 study examining the effect of mint ingestion on GERD. Bulat et al⁷⁹ conducted a randomized double-blind placebo-controlled trial in which the LES tone of healthy volunteers was measured in response to spearmint, and they found no difference in symptoms, sphincter pressure, or reflux episodes with ingestion of spearmint. There were no studies published on the effect of cessation of mint intake on GERD symptoms.

LATE-EVENING MEAL

Postprandial reflux is common in patients with GERD.¹¹⁶ Two studies that evaluated the effect of the timing of the evening meal on 24-hour intragastric acidity in healthy volunteers showed different effects on nocturnal pH. A study⁸⁴ of early (6 PM) vs late (9 PM) dinner demonstrated lower intragastric pH (median pH, 1.39 vs 1.67; $P < .001$) in the later meal setting, but only between midnight and 7 AM. The second study⁸⁵ in 10 healthy patients showed that nocturnal and 24-hour integrated intragastric acidity were unaffected by changes in the timing of the evening meal. In the only study in patients with GERD that examined the effect of late-evening meals, Orr and Harnish⁸⁶ measured esophageal pH and symptoms in 20 patients during a late-evening meal on one night and a meal before 7 PM on another night. There was no difference between the 2 nights in terms of acid exposure, number of reflux episodes, and mean reflux episode duration.⁸⁶ The evidence to routinely recommend avoidance of late-evening meals in patients with GERD remains incomplete.

RECUMBENT POSITION AND HOB ELEVATION

Counseling patients regarding HOB elevation is based on the theory that stomach contents containing acid will more likely reflux into the esophagus while patients are lying flat. Stanciu and Bennett⁸⁰ compared the effect of different body positions (sitting, lying, and elevated HOB) in 63 patients with GERD symptoms using pH monitoring and

showed that compared with patients who slept flat, patients who slept with an elevated HOB (using 28-cm blocks) had significantly fewer reflux episodes, shorter reflux episodes, faster acid clearing, and fewer reflux symptoms. Hamilton et al⁹⁴ conducted a randomized clinical trial comparing sleeping on a wedge, sleep with HOB elevation, and sleeping in the horizontal position and found that sleeping on a wedge was associated with significantly less esophageal acid exposure compared with sleeping in the horizontal position. Sleeping with HOB elevation also showed improved acid exposure, but this difference was not statistically significant.⁹⁴ In another randomized controlled trial⁹⁵ of HOB elevation for 2 weeks in patients with reflux symptoms treated with proton pump inhibitors and cisapride, HOB elevation was not associated with a difference in symptoms or antacid use. The HOB elevation, therefore, seems to be an effective measure to improve the symptoms and physiologic variables in some patients with GERD.

RIGHT LATERAL DECUBITUS POSITION

Several studies have shown that reflux is increased in patients in the right lateral decubitus position. The reason for this phenomenon is not completely clear, but it may be related to increased transient LES relaxations in the right position, or possibly that the gastroesophageal junction lies above the level of gastric acid in the left lateral position.¹¹³ Specifically, total reflux time, average acid clearance, and LES relaxations are significantly prolonged in patients lying on their right side compared with the left lateral decubitus position.⁸¹⁻⁸³ Despite evidence that the right lateral decubitus position can aggravate reflux, the practical aspects of this lifestyle modification make it a challenge to implement in patients with GERD.

CONCLUSIONS

Despite articles in the literature emphasizing the insufficient evidence

to support an association between lifestyle and dietary behaviors and GERD,^{113,117} such interventions continue to be recommended as first-line therapy.^{1,118} Gastroesophageal reflux disease is a chronic disease that affects health-related quality of life.¹¹⁹ The results of this evidence-based approach suggest that although there is physiologic evidence that smoking, alcohol, chocolate, or fatty or citrus food intake may adversely affect symptoms or esophageal pH, there is little evidence that cessation of these agents will improve GERD variables. Elevations in the HOB, left lateral decubitus positioning, and weight loss have been associated with improvement in GERD variables in case-control studies. Larger prospective controlled trials are warranted to conclusively recommend dietary and lifestyle modifications in the treatment of GERD.

Accepted for Publication: November 7, 2005.

Correspondence: Lauren B. Gerson, MD, MSc, Division of Gastroenterology and Hepatology, Stanford University Medical Center, Room A149, Stanford, CA 94305-5202 (lgerson@stanford.edu).

Author Contributions: Dr Gerson had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Financial Disclosure: None.

Funding/Support: This study was supported by a Research Scholar Award from the American Gastroenterological Association (Dr Gerson).

Role of the Sponsor: The American Gastroenterological Association was not involved in the design of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

REFERENCES

1. DeVault KR, Castell DO. Updated guidelines for the diagnosis and treatment of gastroesophageal reflux disease. *Am J Gastroenterol.* 2005; 100:190-200.
2. Gallup Organization. *Heartburn Across America: A Gallup Organization National Survey.* Princeton, NJ: Gallup Organization; 1988.
3. Nebel OT, Fornes MF, Castell DO. Symptomatic gastroesophageal reflux: incidence and precipitating factors. *Am J Dig Dis.* 1976;21:953-956.

4. Dent J, Dodds WJ, Friedman RH, et al. Mechanism of gastroesophageal reflux in recumbent asymptomatic human subjects. *J Clin Invest*. 1980;65:256-267.
5. An evidence-based appraisal of reflux disease management: the Genval Workshop Report. *Gut*. 1999;44(suppl 2):S1-S16.
6. Watanabe Y, Fujiwara Y, Shiba M, et al. Cigarette smoking and alcohol consumption associated with gastro-oesophageal reflux disease in Japanese men. *Scand J Gastroenterol*. 2003;38:807-811.
7. Tibbling L, Gibellino FM, Johansson KE. Is mis-swallowing or smoking a cause of respiratory symptoms in patients with gastroesophageal reflux disease? *Dysphagia*. 1995;10:113-116.
8. Chattopadhyay DK, Greaney MG, Irvin TT. Effect of cigarette smoking on the lower oesophageal sphincter. *Gut*. 1977;18:833-835.
9. Talley NJ, Zinsmeister AR, Schleck CD, Melton LJ III. Smoking, alcohol, and analgesics in dyspepsia and among dyspepsia subgroups: lack of an association in a community. *Gut*. 1994;35:619-624.
10. Nilsson M, Johnsen R, Ye W, Hveem K, Lagergren J. Lifestyle related risk factors in the aetiology of gastro-oesophageal reflux. *Gut*. 2004;53:1730-1735.
11. Kahrilas PJ, Gupta RR. Mechanisms of acid reflux associated with cigarette smoking. *Gut*. 1990;31:4-10.
12. Smit CF, Copper MP, van Leeuwen JA, Schoots IG, Stanojic LD. Effect of cigarette smoking on gastropharyngeal and gastroesophageal reflux. *Ann Otol Rhinol Laryngol*. 2001;110:190-193.
13. Berenson MM, Sontag S, Robinson MG, McCallum RM. Effect of smoking in a controlled study of ranitidine treatment in gastroesophageal reflux disease. *J Clin Gastroenterol*. 1987;9:499-503.
14. Pehl C, Pfeiffer A, Wendl B, Nagy I, Kaess H. Effect of smoking on the results of esophageal pH measurement in clinical routine. *J Clin Gastroenterol*. 1997;25:503-506.
15. Schindlbeck NE, Heinrich C, Dendorfer A, Pace F, Muller-Lissner SA. Influence of smoking and esophageal intubation on esophageal pH-metry. *Gastroenterology*. 1987;92:1994-1997.
16. Wang JH, Luo JY, Dong L, Gong J, Tong M. Epidemiology of gastroesophageal reflux disease: a general population-based study in Xi'an of Northwest China. *World J Gastroenterol*. 2004;10:1647-1651.
17. Stanghellini V. Relationship between upper gastrointestinal symptoms and lifestyle, psychosocial factors and comorbidity in the general population: results from the Domestic/International Gastroenterology Surveillance Study (DIGEST). *Scand J Gastroenterol Suppl*. 1999;231:29-37.
18. Bujanda L. The effects of alcohol consumption upon the gastrointestinal tract. *Am J Gastroenterol*. 2000;95:3374-3382.
19. Kaufman SE, Kaye MD. Induction of gastro-oesophageal reflux by alcohol. *Gut*. 1978;19:336-338.
20. Vitale GC, Cheadle WG, Patel B, Sadek SA, Michel ME, Cuschieri A. The effect of alcohol on nocturnal gastroesophageal reflux. *JAMA*. 1987;258:2077-2079.
21. Rubinstein E, Hauge C, Sommer P, Mortensen T. Oesophageal and gastric potential difference and pH in healthy volunteers following intake of coca-cola, red wine, and alcohol. *Pharmacol Toxicol*. 1993;72:61-65.
22. Grande L, Manterola C, Ros E, Lacima G, Pera C. Effects of red wine on 24-hour esophageal pH and pressures in healthy volunteers. *Dig Dis Sci*. 1997;42:1189-1193.
23. Rosaida MS, Goh KL. Gastro-oesophageal reflux disease, reflux oesophagitis and non-erosive reflux disease in a multiracial Asian population: a prospective, endoscopy based study. *Eur J Gastroenterol Hepatol*. 2004;16:495-501.
24. O'Leary C, McCarthy J, Humphries M, Shananhan F, Quigley E. The prophylactic use of a proton pump inhibitor before food and alcohol. *Aliment Pharmacol Ther*. 2003;17:683-686.
25. Avidan B, Sonnenberg A, Schnell TG, Sontag SJ. No association between gallstones and gastro-oesophageal reflux disease. *Am J Gastroenterol*. 2001;96:2858-2862.
26. Talley NJ, Piper DW. Comparison of the clinical features and illness behaviour of patients presenting with dyspepsia of unknown cause (essential dyspepsia) and organic disease. *Aust N Z J Med*. 1986;16:352-359.
27. Grande L, Monforte R, Ros E, et al. High amplitude contractions in the middle third of the oesophagus: a manometric marker of chronic alcoholism? *Gut*. 1996;38:655-662.
28. Behar J, Sheahan DG, Biancani P, Spiro HM, Storer EH. Medical and surgical management of reflux esophagitis: a 38-month report of a prospective clinical trial. *N Engl J Med*. 1975;293:263-268.
29. O'Brien TF Jr. Lower esophageal sphincter pressure (LESP) and esophageal function in obese humans. *J Clin Gastroenterol*. 1980;2:145-148.
30. Beauchamp G. Gastroesophageal reflux and obesity. *Surg Clin North Am*. 1983;63:869-876.
31. Murray L, Johnston B, Lane A, et al. Relationship between body mass and gastro-oesophageal reflux symptoms: the Bristol Helicobacter Project. *Int J Epidemiol*. 2003;32:645-650.
32. Nilsson M, Johnsen R, Ye W, Hveem K, Lagergren J. Obesity and estrogen as risk factors for gastroesophageal reflux symptoms. *JAMA*. 2003;290:66-72.
33. Diaz-Rubio M, Moreno-Elola-Olaso C, Rey E, Locke GR III, Rodriguez-Artalejo F. Symptoms of gastro-oesophageal reflux: prevalence, severity, duration and associated factors in a Spanish population. *Aliment Pharmacol Ther*. 2004;19:95-105.
34. Nandurkar S, Locke GR III, Fett S, Zinsmeister AR, Cameron AJ, Talley NJ. Relationship between body mass index, diet, exercise and gastro-oesophageal reflux symptoms in a community. *Aliment Pharmacol Ther*. 2004;20:497-505.
35. Locke GR III, Talley NJ, Fett SL, Zinsmeister AR, Melton LJ III. Risk factors associated with symptoms of gastroesophageal reflux. *Am J Med*. 1999;106:642-649.
36. Clements RH, Gonzalez QH, Foster A, et al. Gastrointestinal symptoms are more intense in morbidly obese patients and are improved with laparoscopic Roux-en-Y gastric bypass. *Obes Surg*. 2003;13:610-614.
37. El-Serag HB, Johanson JF. Risk factors for the severity of erosive esophagitis in *Helicobacter pylori*-negative patients with gastroesophageal reflux disease. *Scand J Gastroenterol*. 2002;37:899-904.
38. Foster A, Richards WO, McDowell J, Laws HL, Clements RH. Gastrointestinal symptoms are more intense in morbidly obese patients. *Surg Endosc*. 2003;17:1766-1768.
39. Ruhl CE, Everhart JE. Overweight, but not high dietary fat intake, increases risk of gastroesophageal reflux disease hospitalization: the NHANES I Epidemiologic Followup Study. *Ann Epidemiol*. 1999;9:424-435.
40. Hong D, Khajanchee YS, Pereira N, Lockhart B, Patterson EJ, Swanstrom LL. Manometric abnormalities and gastroesophageal reflux disease in the morbidly obese. *Obes Surg*. 2004;14:744-749.
41. Wajed SA, Streets CG, Bremner CG, DeMeester TR. Elevated body mass disrupts the barrier to gastroesophageal reflux. *Arch Surg*. 2001;136:1014-1018.
42. Fisher BL, Pennathur A, Mutnick JL, Little AG. Obesity correlates with gastroesophageal reflux. *Dig Dis Sci*. 1999;44:2290-2294.
43. Hagen J, Deitel M, Khanna RK, Ilves R. Gastroesophageal reflux in the massively obese. *Int Surg*. 1987;72:1-3.
44. Di Francesco V, Baggio E, Mastromauro M, et al. Obesity and gastro-oesophageal acid reflux: physiopathological mechanisms and role of gastric bariatric surgery. *Obes Surg*. 2004;14:1095-1102.
45. Suter M, Dorta G, Giusti V, Calmes JM. Gastro-oesophageal reflux and esophageal motility disorders in morbidly obese patients. *Obes Surg*. 2004;14:959-966.
46. Lagergren J, Bergstrom R, Nyren O. No relation between body mass and gastro-oesophageal reflux symptoms in a Swedish population based study. *Gut*. 2000;47:26-29.
47. Lundell L, Ruth M, Sandberg N, Bove-Nielsen M. Does massive obesity promote abnormal gastro-oesophageal reflux? *Dig Dis Sci*. 1995;40:1632-1635.
48. Raiha I, Impivaara O, Seppala M, Knuts LR, Sourander L. Determinants of symptoms suggestive of gastroesophageal reflux disease in the elderly. *Scand J Gastroenterol*. 1993;28:1011-1014.
49. Chang CS, Poon SK, Lien HC, Chen GH. The incidence of reflux esophagitis among the Chinese. *Am J Gastroenterol*. 1997;92:668-671.
50. Jaffin BW, Knoepflmacher P, Greenstein R. High prevalence of asymptomatic esophageal motility disorders among morbidly obese patients. *Obes Surg*. 1999;9:390-395.
51. Talley NJ, Howell S, Poulton R. Obesity and chronic gastrointestinal tract symptoms in young adults: a birth cohort study. *Am J Gastroenterol*. 2004;99:1807-1814.
52. Price SF, Smithson KW, Castell DO. Food sensitivity in reflux esophagitis. *Gastroenterology*. 1978;75:240-243.
53. Cohen S, Booth GH Jr. Gastric acid secretion and lower-esophageal-sphincter pressure in response to coffee and caffeine. *N Engl J Med*. 1975;293:897-899.
54. Cohen S. Pathogenesis of coffee-induced gastrointestinal symptoms. *N Engl J Med*. 1980;303:122-124.
55. Boekema PJ, Samsom M, Smout AJ. Effect of coffee on gastro-oesophageal reflux in patients with reflux disease and healthy controls. *Eur J Gastroenterol Hepatol*. 1999;11:1271-1276.
56. DiBaise JK. A randomized, double-blind comparison of two different coffee-roasting processes on development of heartburn and dyspepsia in coffee-sensitive individuals. *Dig Dis Sci*. 2003;48:652-656.
57. Brazer SR, Onken JE, Dalton CB, Smith JW, Schiffman SS. Effect of different coffees on esophageal acid contact time and symptoms in coffee-sensitive subjects. *Physiol Behav*. 1995;57:563-567.
58. Van Deventer G, Kamemoto E, Kuznicki JT, Heckert DC, Schulte MC. Lower esophageal sphincter pressure, acid secretion, and blood gastrin after coffee consumption. *Dig Dis Sci*. 1992;37:558-569.
59. Wendl B, Pfeiffer A, Pehl C, Schmidt T, Kaess H. Effect of decaffeination of coffee or tea on gastro-oesophageal reflux. *Aliment Pharmacol Ther*. 1994;8:283-287.
60. Pehl C, Pfeiffer A, Wendl B, Kaess H. The effect of decaffeination of coffee on gastro-oesophageal reflux in patients with reflux disease. *Aliment Pharmacol Ther*. 1997;11:483-486.
61. Salmon PR, Fedail SS, Wurzner HP, Harvey RF, Read AE. Effect of coffee on human lower oesophageal function. *Digestion*. 1981;21:69-73.
62. Thomas FB, Steinbaugh JT, Fromkes JJ, Mekhjian HS, Caldwell JH. Inhibitory effect of coffee on lower esophageal sphincter pressure. *Gastroenterology*. 1980;79:1262-1266.
63. Wright LE, Castell DO. The adverse effect of chocolate on lower esophageal sphincter pressure. *Am J Dig Dis*. 1975;20:703-707.
64. Murphy DW, Castell DO. Chocolate and heartburn: evidence of increased esophageal acid ex-

- posure after chocolate ingestion. *Am J Gastroenterol.* 1988;83:633-636.
65. Allen ML, Mellow MH, Robinson MG, Orr WC. The effect of raw onions on acid reflux and reflux symptoms. *Am J Gastroenterol.* 1990;85:377-380.
 66. Feldman M, Barnett C. Relationships between the acidity and osmolality of popular beverages and reported postprandial heartburn. *Gastroenterology.* 1995;108:125-131.
 67. Cranley JP, Achkar E, Flesher B. Abnormal lower esophageal sphincter pressure responses in patients with orange juice-induced heartburn. *Am J Gastroenterol.* 1986;81:104-106.
 68. Fass R, Quan SF, O'Connor GT, Ervin A, Iber C. Predictors of heartburn during sleep in a large prospective cohort study. *Chest.* 2005;127:1658-1666.
 69. Crookes P, Hamoui N, Thiesen J, et al. Response of lower esophageal sphincter to ingestion of carbonated beverages. *Gastroenterology.* 1999;116:G0608.
 70. Mangano M, Colombo P, Bianchi PA, Penagini R. Fat and esophageal sensitivity to acid. *Dig Dis Sci.* 2002;47:657-660.
 71. Meyer JH, Lembo A, Elashoff JD, Fass R, Mayer EA. Duodenal fat intensifies the perception of heartburn. *Gut.* 2001;49:624-628.
 72. Holloway RH, Lyrenas E, Ireland A, Dent J. Effect of intraduodenal fat on lower esophageal sphincter function and gastro-oesophageal reflux. *Gut.* 1997;40:449-453.
 73. Nebel OT, Castell DO. Inhibition of the lower esophageal sphincter by fat: a mechanism for fatty food intolerance. *Gut.* 1973;14:270-274.
 74. Becker DJ, Sinclair J, Castell DO, Wu WC. A comparison of high and low fat meals on postprandial esophageal acid exposure. *Am J Gastroenterol.* 1989;84:782-786.
 75. Iwakiri K, Kobayashi M, Kotoyori M, Yamada H, Sugiura T, Nakagawa Y. Relationship between postprandial esophageal acid exposure and meal volume and fat content. *Dig Dis Sci.* 1996;41:926-930.
 76. Pehl C, Waizenhoefer A, Wendl B, Schmidt T, Schepp W, Pfeiffer A. Effect of low and high fat meals on lower esophageal sphincter motility and gastroesophageal reflux in healthy subjects. *Am J Gastroenterol.* 1999;94:1192-1196.
 77. Penagini R, Mangano M, Bianchi PA. Effect of increasing the fat content but not the energy load of a meal on gastro-oesophageal reflux and lower esophageal sphincter motor function. *Gut.* 1998;42:330-333.
 78. Colombo P, Mangano M, Bianchi PA, Penagini R. Effect of calories and fat on postprandial gastro-oesophageal reflux. *Scand J Gastroenterol.* 2002;37:3-5.
 79. Bulat R, Fachnie E, Chauhan U, Chen Y, Tougas G. Lack of effect of spearmint on lower esophageal sphincter function and acid reflux in healthy volunteers. *Aliment Pharmacol Ther.* 1999;13:805-812.
 80. Stanciu C, Bennett JR. Effects of posture on gastro-oesophageal reflux. *Digestion.* 1977;15:104-109.
 81. Katz LC, Just R, Castell DO. Body position affects recumbent postprandial reflux. *J Clin Gastroenterol.* 1994;18:280-283.
 82. van Herwaarden MA, Katzka DA, Smout AJ, Samsom M, Gideon M, Castell DO. Effect of different recumbent positions on postprandial gastroesophageal reflux in normal subjects. *Am J Gastroenterol.* 2000;95:2731-2736.
 83. Khoury RM, Camacho-Lobato L, Katz PO, Mohiuddin MA, Castell DO. Influence of spontaneous sleep positions on nighttime recumbent reflux in patients with gastroesophageal reflux disease. *Am J Gastroenterol.* 1999;94:2069-2073.
 84. Duroux P, Bauerfeind P, Emde C, Koelz HR, Blum AL. Early dinner reduces nocturnal gastric acidity. *Gut.* 1989;30:1063-1067.
 85. Lanzon-Miller S, Pounder RE, McIsaac RL, Wood JR. The timing of the evening meal affects the pattern of 24-hour intragastric acidity. *Aliment Pharmacol Ther.* 1990;4:547-553.
 86. Orr WC, Harnish MJ. Sleep-related gastro-oesophageal reflux: provocation with a late evening meal and treatment with acid suppression. *Aliment Pharmacol Ther.* 1998;12:1033-1038.
 87. Waring JP, Eastwood TF, Austin JM, Sanowski JR. The immediate effects of cessation of cigarette smoking on gastroesophageal reflux. *Am J Gastroenterol.* 1989;84:1076-1078.
 88. Kadakia SC, Kikendall JW, Maydonovitch C, Johnson LF. Effect of cigarette smoking on gastroesophageal reflux measured by 24-h ambulatory esophageal pH monitoring. *Am J Gastroenterol.* 1995;90:1785-1790.
 89. Fraser-Moodie CA, Norton B, Gornall C, Mag-nago S, Weale AR, Holmes GK. Weight loss has an independent beneficial effect on symptoms of gastro-oesophageal reflux in patients who are overweight. *Scand J Gastroenterol.* 1999;34:337-340.
 90. Kjellin A, Ramel S, Rossner S, Thor K. Gastro-oesophageal reflux in obese patients is not reduced by weight reduction. *Scand J Gastroenterol.* 1996;31:1047-1051.
 91. Mathus-Vliegen EM, van Weeren M, van Eerten PV. LOS function and obesity: the impact of untreated obesity, weight loss, and chronic gastric balloon distension. *Digestion.* 2003;68:161-168.
 92. Mathus-Vliegen EM, Tytgat GN. Gastro-oesophageal reflux in obese subjects: influence of overweight, weight loss and chronic gastric balloon distension. *Scand J Gastroenterol.* 2002;37:1246-1252.
 93. Mathus-Vliegen LM, Tytgat GN. Twenty-four-hour pH measurements in morbid obesity: effects of massive overweight, weight loss and gastric distension. *Eur J Gastroenterol Hepatol.* 1996;8:635-640.
 94. Hamilton JW, Boisen RJ, Yamamoto DT, Wagner JL, Reichelderfer M. Sleeping on a wedge diminishes exposure of the esophagus to refluxed acid. *Dig Dis Sci.* 1988;33:518-522.
 95. Pollmann H, Zillesen E, Pohl J, et al. Effect of elevated head position in bed in therapy of gastro-oesophageal reflux [in German]. *Z Gastroenterol.* 1996;34(suppl 2):93-99.
 96. Kahrilas PJ, Gupta RR. The effect of cigarette smoking on salivation and esophageal acid clearance. *J Lab Clin Med.* 1989;114:431-438.
 97. Dua K, Bardan E, Ren J, Sui Z, Shaker R. Effect of chronic and acute cigarette smoking on the pharyngo-upper esophageal sphincter contractile reflex and reflexive pharyngeal swallow. *Gut.* 1998;43:537-541.
 98. Stanciu C, Bennett JR. Smoking and gastro-oesophageal reflux. *BMJ.* 1972;3:793-795.
 99. Mercer CD, Wren SF, DaCosta LR, Beck IT. Lower esophageal sphincter pressure and gastroesophageal pressure gradients in excessively obese patients. *J Med.* 1987;18:135-146.
 100. Stene-Larsen G, Weberg R, Froyshov Larsen I, Bjortuft O, Hoel B, Berstad A. Relationship of overweight to hiatal hernia and reflux oesophagitis. *Scand J Gastroenterol.* 1988;23:427-432.
 101. Wilson LJ, Ma W, Hirschowitz BI. Association of obesity with hiatal hernia and esophagitis. *Am J Gastroenterol.* 1999;94:2840-2844.
 102. Barak N, Ehrenpreis ED, Harrison JR, Sitrin MD. Gastro-oesophageal reflux disease in obesity: pathophysiological and therapeutic considerations. *Obes Rev.* 2002;3:9-15.
 103. Sugerman HJ, DeMaria EJ, Felton WL III, Nakatsuka M, Sismanis A. Increased intra-abdominal pressure and cardiac filling pressures in obesity-associated pseudotumor cerebri. *Neurology.* 1997;49:507-511.
 104. Wisen O, Rossner S, Johansson C. Impaired pancreatico-biliary response to vagal stimulation and to cholecystokinin in human obesity. *Metabolism.* 1988;37:436-441.
 105. Dixon JB, O'Brien PE. Gastroesophageal reflux in obesity: the effect of lap-band placement. *Obes Surg.* 1999;9:527-531.
 106. Angrisani L, Iovino P, Lorenzo M, et al. Treatment of morbid obesity and gastroesophageal reflux with hiatal hernia by Lap-Band. *Obes Surg.* 1999;9:396-398.
 107. O'Brien PE, Dixon JB, Brown W, et al. The laparoscopic adjustable gastric band (Lap-Band): a prospective study of medium-term effects on weight, health and quality of life. *Obes Surg.* 2002;12:652-660.
 108. Smith SC, Edwards CB, Goodman GN. Symptomatic and clinical improvement in morbidly obese patients with gastroesophageal reflux disease following Roux-en-Y gastric bypass. *Obes Surg.* 1997;7:479-484.
 109. Frigg A, Peterli R, Peters T, Ackermann C, Tondelli P. Reduction in co-morbidities 4 years after laparoscopic adjustable gastric banding. *Obes Surg.* 2004;14:216-223.
 110. Perry Y, Courcoulas AP, Fernando HC, Buenaventura PO, McCaughan JS, Luketich JD. Laparoscopic Roux-en-Y gastric bypass for recalcitrant gastroesophageal reflux disease in morbidly obese patients. *JSL.* 2004;8:19-23.
 111. Raftopoulos I, Awais O, Courcoulas AP, Luketich JD. Laparoscopic gastric bypass after antireflux surgery for the treatment of gastroesophageal reflux in morbidly obese patients: initial experience. *Obes Surg.* 2004;14:1373-1380.
 112. Melissas J, Christodoulakis M, Spyridakis M, et al. Disorders associated with clinically severe obesity: significant improvement after surgical weight reduction. *South Med J.* 1998;91:1143-1148.
 113. Meining A, Classen M. The role of diet and lifestyle measures in the pathogenesis and treatment of gastroesophageal reflux disease. *Am J Gastroenterol.* 2000;95:2692-2697.
 114. Penagini R. Fat and gastro-oesophageal reflux disease. *Eur J Gastroenterol Hepatol.* 2000;12:1343-1345.
 115. Feldman M, Friedman LS, Sleisenger MH. *Sleisenger & Fordtran's Gastrointestinal and Liver Disease: Pathophysiology, Diagnosis, Management.* 7th ed. Philadelphia, Pa: WB Saunders Co; 2002.
 116. Johnson LF, Demeester TR. Twenty-four-hour pH monitoring of the distal esophagus: a quantitative measure of gastroesophageal reflux. *Am J Gastroenterol.* 1974;62:325-332.
 117. Holtmann G, Adam B, Liebrechts T. The patient with gastro-oesophageal reflux disease: life-style advice and medication. *Aliment Pharmacol Ther.* 2004;20(suppl 8):24-27.
 118. Kitchin LI, Castell DO. Rationale and efficacy of conservative therapy for gastroesophageal reflux disease. *Arch Intern Med.* 1991;151:448-454.
 119. Gerson LB, Ullah N, Hastie T, Triadafilopoulos G, Goldstein M. Patient-derived health state utilities for gastroesophageal reflux disease. *Am J Gastroenterol.* 2005;100:524-533.